arXiv:1108.1850v1 [math.RA] 9 Aug 2011

DEFINING THE NOTION OF COMPLETE INTERSECTION FOR REGULAR GRADED SKEW CLIFFORD ALGEBRAS

Michaela Vancliff*
Department of Mathematics, P.O. Box 19408,
University of Texas at Arlington,
Arlington, TX 76019-0408
vancliff@uta.edu uta.edu/math/vancliff

Abstract. In recent work of T. Cassidy and the author, a notion of complete intersection was defined for regular skew polynomial rings, defining it using both algebraic and geometric tools. In this article, the definition is modified and extended to a larger class of algebras, namely regular graded skew Clifford algebras. Some of the ideas are also discussed for Auslander-Gorenstein algebras satisfying certain hypotheses.

Introduction

In [2, Definition 3.7], a notion of complete intersection was given for regular skew polynomial rings, defining it using both algebraic and geometric tools. This article addresses whether or not one can extend this definition to a larger class of algebras, namely to regular graded skew Clifford algebras. This latter family of algebras contains the family of regular skew polynomial rings.

Graded skew Clifford algebras were first defined in [2], and may be considered a quantized analog of graded Clifford algebras. They are of interest in the classification of quadratic regular algebras, in part because regular algebras of global dimension at most two are graded skew Clifford algebras, and in part since most quadratic regular algebras of global dimension three may be generated using graded skew Clifford algebras ([2, 5]). Hence, it is likely that

---

1991 *Mathematics Subject Classification.* 16S38, 16E65.

Key words and phrases. complete intersection, regular algebra, Clifford algebra, point module, skew polynomial ring, base point, quadric, Gorenstein.

*M. Vancliff was supported in part by NSF grant DMS-0900239.

graded skew Clifford algebras will be useful in the open problem of classifying quadratic regular algebras of global dimension four. In the spirit of graded Clifford algebras, the results of [2] associate geometric data to graded skew Clifford algebras, and prove that such algebras are quadratic and regular if and only if certain geometric criteria are satisfied (see Theorem 1.4).

In Section 1, we give the definition of a graded skew Clifford algebra and list the main results of [2] that will be useful in our results. In Subsection 1.2, we list the conditions of interest to us in our proposed definition (Definition 2.8) of complete intersection to follow in Section 2; these conditions entail three algebraic tools and one geometric tool. Our results are given in Section 2, with our main results being Theorem 2.2 and Theorem 2.7. These results prove that the algebraic tools of the notion of complete intersection may be extended to regular graded skew Clifford algebras. However, we demonstrate, in Example 2.1, that the geometric tool employed successfully in [2, Definition 3.7] for regular skew polynomial rings fails, in general, on regular graded skew Clifford algebras. Theorem 2.2 also considers similar ideas for Auslander-Gorenstein algebras satisfying certain hypotheses. Such algebras are not usually considered non-commutative analogs of polynomial rings unless they are regular, so it is not clear how useful a notion of complete intersection would be for such algebras if they are not regular.

1. Graded Skew Clifford Algebras

In Subsection 1.1, we define graded skew Clifford algebras and list the results of [2] that will be of interest in Section 2. We list, in Subsection 1.2, some conditions that will be useful in our proposed definition (Definition 2.8) of complete intersection in Section 2.

Throughout the article, $\Bbbk$ denotes an algebraically closed field with $\mathrm{char}(\Bbbk) \neq 2$, $M(n,\ \Bbbk)$ denotes the space of $n \times n$ matrices with entries in $\Bbbk$, and $\Bbbk^\times = \Bbbk \setminus \{0\}$.

1.1. **The Algebras and Known Results.**

The following notion of symmetry for square scalar matrices generalizes the standard notions of symmetry and skew-symmetry for such matrices.

Definition 1.1. [2] For $\{i, j\} \subset \{1, \ldots, n\}$, let $\mu_{ij} \in \Bbbk$ satisfy $\mu_{ij}\mu_{ji} = 1$ for all $i \neq j$, and write $\mu = (\mu_{ij}) \in M(n, \Bbbk)$. A matrix $M \in M(n, \Bbbk)$ is called *μ-symmetric* if $M_{ij} = \mu_{ij} M_{ji}$ for all $i, j = 1, \ldots, n$.

Definition 1.2. [2] With μ as above, suppose $\mu_{ii} = 1$ for all i, and fix μ-symmetric matrices $M_1, \ldots, M_n \in M(n, \Bbbk)$. A *graded skew Clifford algebra* associated to μ and $M_1, \ldots, M_n$ is a graded $\Bbbk$-algebra on degree-one generators $x_1, \ldots, x_n$ and on degree-two generators $y_1, \ldots, y_n$ with defining relations given by:

(a) $x_i x_j + \mu_{ij} x_j x_i = \sum_{k=1}^{n} (M_k)_{ij} y_k$ for all $i, j = 1, \ldots, n$, and
(b) the existence of a normalizing sequence $\{r_1, \ldots, r_n\}$ of homogeneous elements of degree two that span $\Bbbk y_1 + \cdots + \Bbbk y_n$.

The reader should note that if $\mu_{ij} = 1$ for all i, j, and if y_k is central in the algebra for all k, then the algebra is a graded Clifford algebra. On the other hand, taking $M_1, \ldots, M_n$ to be diagonal and linearly independent shows that regular skew polynomial rings on finitely-many generators are graded skew Clifford algebras.

Although graded skew Clifford algebras need not be quadratic nor regular in general, a simple geometric criterion was established in [2, Theorem 4.2] for determining when such an algebra is quadratic and regular. In order to state that result, we first define a (non-commutative) quadric system.

Definition 1.3. [2] Fix μ as in Definition 1.2, and let S denote the regular skew polynomial ring

$$\frac{T}{\langle z_j z_i - \mu_{ij} z_i z_j \ : \ i, j = 1, \ldots, n \ \rangle},$$

where T is the tensor (free) algebra on generators $z_1, \ldots, z_n$. We may identify $\mathbb{P}(S_1^*)$ with $\mathbb{P}^{n-1}$, and, under this identification, we denote the zero locus, in $\mathbb{P}^{n-1} \times \mathbb{P}^{n-1}$, of the defining relations of S by Z. For any $q \in S_2$, we call the zero locus of q in Z the *quadric associated to* q, and denote it by $\mathcal{V}_S(q)$; in other words, $\mathcal{V}_S(q) = \mathcal{V}(\hat{q}) \cap Z$, where $\hat{q}$ is any lift of q to T_2, and $\mathcal{V}(\hat{q})$ is the zero locus of $\hat{q}$ in $\mathbb{P}^{n-1} \times \mathbb{P}^{n-1}$. The span of elements $q_1, \ldots, q_m$ in S_2 will be called the *quadric system* associated to $q_1, \ldots, q_m$, and if $\mathcal{V}_S(q_1) \cap \cdots \cap \mathcal{V}_S(q_m)$ is empty, then the quadric system is said to be *base-point free*. If a quadric system is given by a normalizing sequence in S, then it is called a *normalizing quadric system.*

Given μ and S as above, and a μ-symmetric matrix M, one may associate to M an element $q \in S_2$ as follows:

$$q = \begin{bmatrix} z_1 & \dots & z_n \end{bmatrix} M \begin{bmatrix} z_1 \\ \vdots \\ z_n \end{bmatrix} \in S_2.$$

Theorem 1.4. [2] *A graded skew Clifford algebra $A = A(\mu, M_1, \dots, M_n)$ as in Definition 1.2 is a quadratic, Auslander-regular algebra of global dimension n that satisfies the Cohen-Macaulay property with Hilbert series $1/(1-t)^n$ if and only if the quadric system associated to $M_1, \dots, M_n$ is normalizing and base-point free; in this case, A is a noetherian regular domain and is unique up to isomorphism.*

The "base-point free" property in the previous result is referring to the associated quadric system in the skew polynomial ring S yielding a complete intersection in the sense of Definition 1.6 below.

Example 1.5. Let $n = 3$ and $\mu_{ij} = 1$ for all i, j, so, in this example, S is the polynomial ring on three variables and all sequences in S are normalizing. Taking

$$M_1 = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad M_2 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad M_3 = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 2 \end{bmatrix},$$

we obtain the quadric system

$$z_1^2, \quad z_2^2, \quad z_1 z_2 + z_3^2,$$

which is base-point free. By the preceding result, the graded (skew) Clifford algebra associated to μ and the M_k is the quadratic regular algebra

$$\frac{\Bbbk\langle x_1,\ x_2,\ x_3\rangle}{\langle x_1x_2 + x_2x_1 - x_3^2,\ x_1x_3 + x_3x_1,\ x_2x_3 + x_3x_2\rangle}.$$

We will return to this example in Section 2.

1.2. **The Conditions.**

In order to address the notion of complete intersection for a non-commutative graded algebra A on n generators with a normalizing sequence $F = \{f_1, \dots, f_n\} \subset A$ of homogeneous elements of positive degree, we will be interested in four (possibly equivalent) conditions on A and F as follows:

I. F is a regular sequence in A;

II. $\dim_{\Bbbk}(A/\langle F\rangle) < \infty$;

III. for each $k = 1, \ldots, n$, $\operatorname{GKdim}(A/\langle f_1, \ldots, f_k\rangle) = n - k$;

IV. there are no right point modules M over A such that $Mf_k = 0$ for all k.

If $\deg(f_k) = 2$ for all k, and if the right point modules over A are parametrized by the zero locus of the degree-2 defining relations of A in $\mathbb{P}(A_1^*) \times \mathbb{P}(A_1^*)$, then condition IV is equivalent to the normalizing quadric system $\{f_1, \ldots, f_n\}$ being base-point free.

The definition of complete intersection in [2] that we wish to extend to regular graded skew Clifford algebras is based on [2, Corollary 3.6], which proves that conditions I-IV are equivalent in the case that A is a regular skew polynomial ring. This motivates the following definition in [2].

Definition 1.6. [2] Let A denote a regular skew polynomial ring on n generators. If F is a normalizing sequence in A of n homogeneous elements of positive degree, then we call $A/\langle F\rangle$ a *complete intersection* if the equivalent conditions I-IV hold.

In Definition 2.8, we extend (with some modification) Definition 1.6 to regular graded skew Clifford algebras.

2. Complete Intersections

It is a natural problem to seek the algebras A for which two or more of conditions I-IV from Subsection 1.2 are equivalent. In this section, we show that if A is a regular graded skew Clifford algebra, then conditions I-III are equivalent, but that, in general, condition IV is not equivalent to conditions I-III. We first address this latter issue via an example.

Example 2.1. Let A denote the regular graded (skew) Clifford algebra given in Example 1.5. Since A is a graded Clifford algebra, the elements x_1^2, x_2^2 and x_3^2 are central in A. It follows that the sequence $F = \{f_1 = x_1^4,\ f_2 = x_2^4,\ f_3 = x_2^2x_1^2\}$ is central (and so normalizing) in A. Clearly, F is not a regular sequence, since A is a domain, the elements in F are all homogeneous of the same degree, and $x_2^2 f_3 = f_2 x_1^2$. Using Bergman's Diamond Lemma, it is straightforward to see that $\mathrm{GKdim}(A/\langle F\rangle) = 1$, from which we obtain that conditions I-III all fail. However, the right point modules over A are given by points in $\mathbb{P}^2 \times \mathbb{P}^2$ of the form:

$$((a,\ b,\ 0),\ (a,\ -b,\ 0)) \quad \text{and} \quad ((2a^2,\ -b^2,\ 2ab),\ (2a^2,\ -b^2,\ -2ab))$$

for all $(a,\ b) \in \mathbb{P}^1$. Therefore, to find the right point modules over A on which the f_k vanish, we apply each f_k in turn to points of the form

$$((a,\ b,\ 0),\ (a,\ -b,\ 0),\ (a,\ b,\ 0),\ (a,\ -b,\ 0)) \in (\mathbb{P}^2)^4$$

and

$$((2a^2,\ -b^2,\ 2ab),\ (2a^2,\ -b^2,\ -2ab),\ (2a^2,\ -b^2,\ 2ab),\ (2a^2,\ -b^2,\ -2ab)) \in (\mathbb{P}^2)^4,$$

where $(a,\ b) \in \mathbb{P}^1$. Since this computation yields no solution, condition IV holds. Hence, for this algebra A and the given sequence F, condition IV is not equivalent to conditions I-III.

In spite of the preceding example, the next result shows that conditions I-IV are equivalent for some normalizing sequences in certain graded algebras; the technical hypotheses are motivated by results in [4].

Theorem 2.2. *Let $A = \oplus_{i\geq 0} A_i$ denote an $\mathbb{N}$-graded, Auslander-Gorenstein $\Bbbk$-algebra of finite injective dimension with $\dim_{\Bbbk}(A_0) < \infty$ that satisfies the Cohen-Macaulay property with GKdim$(A) = n \in \mathbb{N} \setminus \{0\}$. Suppose F is a normalizing sequence in A of n homogeneous elements of positive degree.*

(a) *If A is noetherian, then conditions I-III are equivalent.*

(b) *If $A/\langle F\rangle \cong B/\langle G\rangle$, where B is a skew polynomial ring on n generators and G is a normalizing sequence in B of n homogeneous elements of positive degree, then conditions I-IV are equivalent for A and F.*

Proof. (a) Part (a) is [2, Theorem 2.5]. (We remark that the proof of [2, Theorem 2.5] uses results from [4] that assume $\dim_{\Bbbk}(A_0) < \infty$, although this is not explicitly stated in [2, Theorem 2.5].)

(b) Suppose $A/\langle F\rangle \cong B/\langle G\rangle$, where B and G are as given in the statement. It follows that $A/\langle F\rangle$ is noetherian, so [1, Lemma 8.2], implies that A is noetherian. Hence, conditions I-III are equivalent by (a).

Condition IV holds if and only if $A/\langle F\rangle$ has no right point modules, and so it holds if and only if there are no right point modules N over B such that $Ng = 0$ for all $g \in G$. By [2, Corollary 3.6], this situation happens if and only if $\dim(B/\langle G\rangle) < \infty$. It follows that condition IV is equivalent to condition II. ■

Since many known Auslander-regular (respectively, regular) algebras satisfy the hypotheses of Theorem 2.2, we have that Theorem 2.2 may be applied to many such algebras. However, algebras satisfying the hypotheses of Theorem 2.2 that are not regular are not usually considered non-commutative analogs of polynomial rings, so it is unclear how useful a notion of complete intersection would be for such algebras.

In Examples 2.3 and 2.4, we present applications of Theorem 2.2. The reader should note that Example 2.4 is not a graded skew Clifford algebra.

Example 2.3. Let A denote the regular graded Clifford algebra given in Example 1.5, which satisfies the hypotheses of Theorem 2.2. Let $f_i = x_i^2$ for all i. Since $\{f_1,\ f_2\}$ is base-point free, one might first think the geometric data is not helpful for determining information about the full sequence $F = \{f_1,\ f_2,\ f_3\}$. However, $A/\langle F\rangle \cong B/\langle G\rangle$, where

$$B = \frac{\Bbbk\langle x_1,\ x_2,\ x_3\rangle}{\langle x_1x_2 + x_2x_1,\ x_1x_3 + x_3x_1,\ x_2x_3 + x_3x_2\rangle},$$

and $G = \{x_1^2,\ x_2^2,\ x_3^2\} \subset B$. By Theorem 2.2(b), conditions I-IV are equivalent for A and F. Since F is base-point free, condition IV holds, so conditions I-III also hold for A and F.

Example 2.4. Let A denote the coordinate ring of quantum 2×2 matrices, which is the quadratic $\Bbbk$-algebra on generators a, b, c, d with defining relations

$$\begin{aligned} ab &= qba, & bd &= qdb,\\ ac &= qca, & cd &= qdc,\\ bc &= cb, & ad - da &= (q - q^{-1})bc, \end{aligned}$$

where $q \in \Bbbk^\times$ and $q^2 \neq 1$ ([3]). By [7], this algebra is an Auslander-regular algebra of global dimension four with Hilbert series $1/(1-t)^4$ and it satisfies the hypotheses of Theorem 2.2. However, it is not a graded skew Clifford algebra. Let $F = \{bc,\ b^2,\ c^2,\ ad\}$, which is a normalizing nonregular sequence in A, so condition I fails. However, $A/\langle F\rangle \cong B/\langle G\rangle$, where

B is the $\Bbbk$-algebra on generators a, b, c, d with defining relations

$$\begin{aligned} ab &= qba, & bd &= qdb, \\ ac &= qca, & cd &= qdc, \\ bc &= cb, & ad &= da, \end{aligned}$$

and $G = \{bc,\ b^2,\ c^2,\ ad\} \subset B$. By Theorem 2.2, conditions I-IV are equivalent for A and F, so none of the conditions hold.

In the next result, it should be noted that it is not assumed that the quadric system associated to the graded skew Clifford algebra A is linearly independent, nor that A is quadratic (until (d)).

Proposition 2.5. *Let A denote a graded skew Clifford algebra and let F denote the normalizing sequence of length n given in Definition 1.2(b).*

- (a) *The algebra A is noetherian.*
- (b) *Conditions II and IV hold for A and F.*
- (c) *If F is regular in A, then A satisfies the hypotheses of Theorem 2.2 and has Hilbert series $H(t) = 1/(1-t)^n$, and conditions I-IV hold for A and F.*
- (d) *If F is regular in A and if A is quadratic, then A is Auslander-regular of global dimension n and the quadric system associated to A is normalizing and base-point free.*

Proof. In the notation of Definition 1.2, we have

$$\frac{A}{\langle F \rangle} \cong \frac{A}{\langle \sum \Bbbk y_k \rangle} \cong \frac{\Bbbk\langle x_1, \ldots, x_n \rangle}{\langle x_i x_j + \mu_{ij} x_j x_i : 1 \leq i,\ j \leq n \rangle} \cong \frac{B}{\langle G \rangle},$$

where B is a regular skew polynomial ring on generators $x_1, \ldots, x_n \in B_1$ and $G = \{x_1^2, \ldots, x_n^2\}$.

(a) Since $B/\langle G \rangle$ is noetherian, it follows that A is noetherian by [1, Lemma 8.2].

(b) The algebra $B/\langle G \rangle$ has finite dimension and has no right point modules. Hence, conditions II and IV hold for A and F.

(c) The sequence G is normalizing in B and is base-point free. By [2, Corollary 3.6], G is a regular sequence in B, and, by [4, Theorem 5.10], $B/\langle G \rangle$ is Auslander-Gorenstein of finite injective dimension and satisfies the Cohen-Macaulay property. Thus, if F is regular in A, then, by [4, Theorem 5.10], A satisfies the hypotheses of Theorem 2.2, so conditions I-IV hold for A and F. Since the Hilbert series of $B/\langle G \rangle$ is $h(t) = (1+t)^n$ and since F is regular and normalizing in A, the Hilbert series of A is $H(t) = 1/(1-t)^n$.

(d) By [6, Theorem 1.5], $B/\langle G\rangle$ is Koszul, since B is Koszul and G is regular and normalizing. Thus, $A/\langle F\rangle$ is Koszul. Since A is quadratic and F is regular and normalizing, A is Koszul by [6, Theorem 1.5]. By (c), the Koszul dual of A has Hilbert series $h(t) = (1+t)^n$. It follows that the right trivial module $\Bbbk_A$ has projective dimension n, so A has global dimension n. The result follows by Theorem 1.4. ■

The following example shows that A being quadratic does not follow from F being regular in Proposition 2.5(c)(d).

Example 2.6. Let A denote the graded Clifford algebra given by the symmetric matrices

$$M_1 = \begin{bmatrix} 2 & -1 \\ -1 & 0 \end{bmatrix}, \quad M_2 = \begin{bmatrix} 0 & -1 \\ -1 & 2 \end{bmatrix}.$$

Hence, A is a factor algebra of the $\Bbbk$-algebra A' on degree-1 generators $x,\ y$ with defining relations

$$xy + yx = -x^2 - y^2, \qquad x^2y = yx^2,$$

and the associated quadric system is given by $z_1(z_1 - z_2)$, $z_2(z_1 - z_2)$, which has base-point $(1,\ 1)$. Since x^2 and y^2 are central elements in A', we have that $A = A'$. Since a basis for A_m is $\{x^iy^j : i + j = m\}$, for all $m \in \mathbb{N}$, the Hilbert series of A is $H(t) = 1/(1-t)^2$, and, since $A/\langle x^2, y^2\rangle$ has Hilbert series $h(t) = (1+t)^2$, it follows that $\{x^2, y^2\}$ is a regular normalizing sequence in A. Thus, by Proposition 2.5(c), conditions I-IV hold for A and the sequence $F = \{x^2, y^2\}$. Although A satisfies the hypotheses of Theorem 2.2, one should note that A is not quadratic and not regular (since $(x+y)^2 = 0$ in A). The author thanks James Zhang of the University of Washington for providing this example.

Theorem 2.7. *Let A denote a regular graded skew Clifford algebra of global dimension n in the sense of Theorem 1.4.*

(a) *The algebra A satisfies the hypotheses of Theorem 2.2.*
(b) *Conditions I-IV hold for A and the normalizing sequence from Definition 1.2(b).*
(c) *If F is any normalizing sequence in A of n homogeneous elements of positive degree, then conditions I-III are equivalent.*

Proof. The hypotheses imply that the normalizing sequence from Definition 1.2(b) is regular. Hence, (a) and (b) follow from applying Proposition 2.5(c). By Proposition 2.5(a), A is noetherian, so (c) follows from Theorem 2.2(a). ■

Given Theorem 2.7(c), we propose the following definition.

Definition 2.8. Let A denote a regular graded skew Clifford algebra of global dimension n in the sense of Theorem 1.4. If F is a normalizing sequence in A of n homogeneous elements of positive degree, then we call $A/\langle F\rangle$ a *complete intersection* if the equivalent conditions I-III hold.

Question 2.9.

(a) Is it possible to define the notion of complete intersection in the spirit of Definition 2.8 to algebras more general than regular graded skew Clifford algebras? Moreover, are any of the hypotheses of Theorem 2.2 unnecessary?

(b) Would condition IV be more widely applicable if it were to use truncated point modules of appropriate lengths in place of point modules? In order to have a geometric condition be equivalent to conditions I-III, is a situation similar to that of Theorem 2.2(b) necessary?

(c) In Example 2.1, a regular graded skew Clifford algebra and sequence were given for which condition IV was not equivalent to conditions I-III. Does a similar example exist where the sequence contains only homogeneous elements of degree two?

References

[1] M. Artin, J. Tate and M. Van den Bergh, Some Algebras Associated to Automorphisms of Elliptic Curves, *The Grothendieck Festschrift* **1**, 33-85, Eds. P. Cartier et al., Birkhäuser (1990).

[2] T. Cassidy and M. Vancliff, Generalizations of Graded Clifford Algebras and of Complete Intersections, *J. Lond. Math. Soc.* **81** (2010), 91-112.

[3] L. D. Faddeev, N. Yu. Reshetikhin and L. A. Takhtadzhyan, Quantization of Lie Groups and Lie Algebras, *Leningrad Math. J.* **1** No. 1 (1990), 193-225.

[4] T. Levasseur, Some Properties of Non-commutative Regular Graded Rings, *Glasgow Math. J.* **34** (1992), 277-300.

[5] M. Nafari, M. Vancliff and Jun Zhang, Classifying Quadratic Quantum $\mathbb{P}^2$s by using Graded Skew Clifford Algebras, revised preprint, June 2011.

[6] B. Shelton and C. Tingey, On Koszul Algebras and a New Construction of Artin-Schelter Regular Algebras, *J. Algebra* **241** No. 2 (2001), 789-798.

[7] M. Vancliff, Quadratic Algebras Associated with the Union of a Quadric and a Line in $\mathbb{P}^3$, *J. Algebra* **165** No. 1 (1994), 63-90.